arXiv:1210.8060v1 [cond-mat.stat-mech] 30 Oct 2012

Combinatorics of lattice animals by simple group–theoretic approach

N. Haug[1,2] and S. Nechaev[1,3]
[1] *LPTMS, Université de Paris Sud 11, 91405 Orsay Cedex, France*
[2] *Institut für Physik, Humboldt-Universität zu Berlin, 12489 Berlin, Germany*
[3] *P.N. Lebedev Physical Institute of the Russian Academy of Sciences, 119991 Moscow, Russia*

Using the commutation relations in the locally free group we give a simple combinatoric derivation of the partition function of (1+1)–dimensional directed lattice animals with a fixed set of roots on a strip. Comparing our approach to the two–dimensional hard square model, we establish connections between the group–theoretic and the algebraic methods of the solution of some two–dimensional lattice spin systems.

PACS numbers:

I. INTRODUCTION

The term "directed lattice animals" is used as a collective name for several related models describing the growth of aggregates, as the growth of molecular layers on substrates. Directed animals in the two–dimensional geometry are structures of occupied and unoccupied nodes on a lattice strip of arbitrary length along one dimension and with a finite width along the other one. It is convenient to align the lattice diagonally to the strip's edges. The occupied sites on the lowest row are the roots (the "source points") and the animal has to satisfy the condition that each occupied site can be reached from at least one root over a path which does only contain occupied sites and which never goes opposite to the preferred direction. If we ascendingly label the nodes in the lowest row from left to right, the set of roots is given by the indices of the occupied nodes [7]. Figure 1 shows an example.



Figure 1: Directed animal on a strip of width $n = 3$ with set of roots $C = \{1, 2\}$. Occupied nodes are drawn black, the roots are highlighted grey. The vertical arrow indicates the preferred direction.

In this work, we give a simple derivation of the partition function of directed lattice animals with fixed root configuration on a strip of finite width. This problem has been solved exactly in [2] by the diagonalization of a specially designed transfer matrix via algebraic methods borrowed from integrable systems. Specifically, in [2], the authors defined the transfer matrix T between row configurations represented in a spin space and computed the eigenvectors of T by finding the appropriate scattering R–matrix as adopted in the theory of integrable systems. The result shows an exponential growth proportional to 3^s in the limit of large numbers of occupied nodes s.

The approach exploited in our work uses the correspondence between directed animals and heaps of pieces. These heaps have been introduced in [5] as a geometric interpretation of the Cartier–Foata's partially commutative monoid. The combinatorial solution described in our paper relies on the natural commutative structure of the "locally free" group [4], which describes available configurations of directed animals on a given lattice. The main idea consists of establishing a bijection between word counting in the locally free group and the particular configuration of the directed animal. The most involved part of the method is the correct account of the specific root configurations.

By presenting our method we would like to direct attention towards possible hidden connections between the algebraic method developed in [2] and our geometric group–theoretic approach based on the concept of the locally free group.

II. THE MAIN RESULT

We compute the partition function of directed animals with fixed roots by using the correspondence between lattice animals and a partially commutative semi–group, more precisely the locally free idempotent semi–group. We prove the following theorem:

Theorem 1. *The number $\mathcal{N}_n(C,s)$ of directed animals on a strip of width n with s occupied sites and set of roots $C = \{k_1, \dots, k_\ell\}$ with $k_1 < k_2 < \dots < k_\ell$ is given by the expression*

$$\mathcal{N}_n(C,s) \;=\; \sum_{m=1}^{4n+3} \frac{(-1)^{m+1}}{2n+2} \sin\frac{m\pi}{n+1} \left(\sin\frac{m\pi}{2n+2}\right)^{-\ell} \prod_{i=0}^{\ell} \left(\sin\frac{(2h_i+1)m\pi}{2n+2}\right) \left(1+2\cos\frac{m\pi}{n+1}\right)^{s-1} \qquad (1)$$

where $h_0 = k_1 - 1$*,* $h_i = k_{i+1} - k_i - 1$ *for* $1 \leq i < l$ *and* $h_\ell = n - k_\ell + \frac{1}{2}$*.*

This paper is organized as follows. In section III, we recall the correspondence between directed animals and heaps of pieces and give the definition of the locally free group. In section IV, we consider lattice animals with a single root before generalizing to arbitrary numbers of roots in section V. Finally, we summarize our results in section VI.

III. GROUP–THEORETIC FORMULATION OF THE PROBLEM

A lattice animal as shown in figure 1 can equivalently be drawn as a heap of 2–dimensional rectangular boxes which have discrete horizontal positions (columns) and which can only touch each other at their corners. The roots of the heap are the boxes which stand on the bottom of a column. Figure 2 shows the equivalence of the two pictures.

We write $\omega_n(\{C\}, s)$ for the number of heaps of s boxes on n columns with a set of roots C. As can be deduced from figure 2, its correspondence to the partition functions of directed animals is

$$\mathcal{N}_n(\{k_1,\dots\},s) \;=\; \omega_{2n}(\{2k_1-1,\dots\},s) \qquad (2)$$



Figure 2: A lattice animal on a strip of width 3 drawn as a heap of boxes on 6 columns.

The combinatorial treatment of heaps of pieces was formalized by G. Viennot [5]. The boxes which we consider (called dimers in [5]) are a special kind of pieces. Viennot showed that each heap of pieces can be seen as the graphical representation of an element of a partially commutative semi–group. For our heaps of boxes, this is the *locally free idempotent* semi–group.

Definition 1 (Locally free idempotent semi–group)**.** *A semi–group of rank* n *with the generating set* $\{f_1,\dots,f_n\}$ *is called a locally free idempotent semi–group* $\mathcal{LFI}^+_{n+1}$ *if it satisfies following relations.*

$$f_i f_j \;=\; f_j f_i \quad \textit{for} \;\; |i-j| \geq 2 \qquad (3a)$$

$$(f_i)^2 \;=\; f_i \quad \textit{for} \;\; i = 1\dots n \qquad (3b)$$

and if each two generators $f_i, f_{i+1},\;\; i = 1\dots n{-}1$ *produce a free sub–semi–group of* $\mathcal{LFI}^+_{n+1}$*.*

We will always refer to the generating set specified in the definition above. Each of the generators corresponds to the act of dropping a box in one of the columns. Figure 3 shows the meaning of the commutation relations in the heaps picture.



Figure 3: The uppermost picture shows that the dropping of boxes on adjacent columns does not commute, whereas it commutes if there is at least one column inbetween (middle). The sequential dropping of two boxes in the same column is the same as dropping only one box (below).

Each element of $\mathcal{LFI}^+_{n+1}$ can be written as a product of the generators like

$$g = f_{\alpha_1} \cdot f_{\alpha_2} \cdots f_{\alpha_s} \qquad (4)$$

Here, s can be any natural number. Each such product can be written as a *word*

$$W = (\alpha_1, \alpha_2, \dots, \alpha_s) \qquad (5)$$

However, due to the partial commutativity of the generators, the word representation of a semi–group element is in general not unique. Two words which represent the same semi–group element belong to the same equivalence class. For example, it is

$$(1,3,2) \sim (3,1,2) \tag{6}$$

In order to find a unique word representation of each element of $\mathcal{LFI}^+_{n+1}$, the authors of [4] introduced the *ordered word form.*

Definition 2. *An s–tuple $W = (\alpha_1, \ldots, \alpha_s)$ with $\alpha_i \in \{1, \ldots, n\}$ for $i = 1 \ldots s$ has the ordered form, if and only if it satisfies the following conditions.*

$$\begin{array}{llll} \textit{if} & \alpha_i = 1 & \textit{then} & \alpha_{i+1} \in \{2, \ldots, n\} \qquad \text{(i)} \\ \textit{if} & 1 < \alpha_i < n & \textit{then} & \alpha_{i+1} \in \\ & & & \{\alpha_i - 1, \alpha_i + 1, \ldots, n\} \ \text{(ii)} \\ \textit{if} & \alpha_i = n & \textit{then} & \alpha_{i+1} = n - 1 \qquad \text{(iii)} \end{array}$$

The conditions (i)-(iii) are visualized in figure 4.



Figure 4: The arrows point to the values the $(i+1)$th letter of the ordered word can take if the ith letter is x.

Each element of $\mathcal{LFI}^+_{n+1}$ is represented in one and only one way as an ordered word. The formal expression for the number $\theta_n(x, s)$ of words of length s with first letter x in the box of size n is

$$\theta_n(x, s) = \#\{\,(x, \alpha_2, \ldots, \alpha_s) \mid \text{(i)-(iii)}\,\} \tag{7}$$

The function $\theta_n(x, s)$ satisfies the recursion relation which straightforwardly follows from the conditions (i)-(iii) and figure 4:

$$\begin{cases} \theta_n(x, s+1) = \theta_n(x-1, s) + \sum_{y=x+1}^{n} \theta_n(y, s), \\ \theta_n(x, 1) = \delta_{x,x_0}, \\ \theta_n(0, s) = \theta(n+1, s) = 0. \end{cases} \tag{8}$$

The generating function

$$\Theta_n(x, \lambda) = \sum_{s=1}^{\infty} \theta_n(x, s) \lambda^{-s}$$

can be easily computed by solving (8) (see [4] for details). It reads

$$\Theta_n(x, \lambda) = (\lambda + 1)^{\frac{n-x}{2}} \frac{\sin \phi n}{\sin \phi (n+2)} \tag{9}$$

In particular,

$$\Theta_n(n, \lambda) = \frac{\sin \phi n}{\sin \phi (n+2)} \tag{10}$$

Here and in the following, the angle ϕ is defined via the identity

$$\phi = \arccos \frac{\sqrt{\lambda + 1}}{2} \tag{11}$$

In eqs. (9)–(10) $\lambda \in [-1, 3]$ and all poles lie within this interval [8].

In order to derive results for the lattice animal problem it is necessary to find out how a given root translates to the corresponding word representation. This rather nontrivial task will be adressed in the next two sections.

IV. HEAPS WITH A SINGLE ROOT

In this chapter we search for the number of heaps with a single root at a given site. The question we ask can be rephrased as follows: “Which condition (additional to conditions (i)-(iii)) does the word representing a heap with one single root satisfy?”

Suppose we want to generate a heap with a single root in the kth column. Obviously, the first letter of the word has to be k. Since we do not want that the second box falls on the ground, we have to make sure that it falls ontop of the first one. Consequently, α_2 has to be either $k-1$ or $k+1$. In order to prevent that α_2 takes any other value than one of these two, we put a “wall” at site $k+2$ (see figure 5). As soon as the first box is dropped in column $k+1$, we can shift the wall to the right by one step, because then, any box dropped in the $k+2$th column will fall ontop of the one in column $k+1$. When the first box is dropped in column $k+2$, we can shift our wall again, and so on.



Figure 5: Reduced set of possible values of α_2 compatible with the ordered word form and preventing of having roots other than in position $\alpha_1 = x$. Jumps from position $\alpha_1 = x$ to positions $\alpha_2 = x+2, ..., n$ are blocked.

Algebraically, the condition which ensures that the ordered word represents a heap *with a single root* located in the column x (see figure 5), reads

$$\alpha_j < \max\{\,\alpha_i\,|\,1 \le j < i\,\} + 2 \quad \forall\, j \in \{2,\ldots,s\} \qquad \text{(iv)}$$

With (iv), we can now write the partition function of heaps in n columns with s elements and a single root in the xth column (see figure 5) as

$$\omega_n(\{x\},s) = \#\{\,(x,\ldots,\alpha_s)\,|\,\text{(i)-(iv)}\,\} \tag{12}$$

Condition (iv) is automatically satisfied for $\alpha_1 = n$. Consequently it is

$$\omega_n(\{n\},s) = \theta_n(n,s) \tag{13}$$

With the use of equation (13), the following theorem can be derived.

Theorem 2. *The partition function $\omega_n(\{k\},s)$ of heaps with a single root at site k satisfies the equation*

$$\omega_n(\{k\},s) = \theta_k(k,s) + \sum_{s_0+s_1=s} \theta_k(k,s_0)\,\theta_{k+1}(k+1,s_1) + \cdots + \sum_{s_0+\cdots+s_{n-k}=s} \theta_k(k,s_0)\ldots\theta_n(n,s_{n-k}) \tag{14}$$

Since the proof is straightforward, it is not carried out here.

The series (14) has the form of a convolution, so we step over to the generating function

$$\Omega_n(\{k\},\lambda) = \sum_{s=1}^{\infty} \omega_n(\{k\},s)\,\lambda^{-s} \tag{15}$$

for which equation (14) transforms to

$$\Omega_n(\{k\},\lambda) = \sum_{d=0}^{n-k} \prod_{j=0}^{d} \Theta_{k+j}(k+j,\lambda) \tag{16}$$

By inserting (9), we get after some algebra

$$\Omega_n(\{k\},\lambda) = \frac{\sin k\phi\ \sin(n-k+1)\phi}{\sin\phi\ \sin(n+2)\phi} \tag{17}$$

for $\lambda \in [-1,3]$.

V. GENERALIZATION TO ARBITRARY NUMBERS OF ROOTS

Now we are in the position to generalize our results to arbitrary numbers of roots.

Suppose that a heap of s pieces possesses two roots in k_1 and k_2 and that $k_1 < k_2$. The representing word has the form

$$W = (k_1, \alpha_2, \ldots, \alpha_{s_1}, k_2, \alpha_{s_1+2}, \ldots, \alpha_{s_1+s_2}) \tag{18}$$

with $s_1 + s_2 = s$ and $\alpha_i \neq k_2\ \forall\ i = 1\ldots s_1$. Since the first part of W, $(k_1, \alpha_2, \ldots, \alpha_{s_1})$, does not generate a root except in k_1, it has to satisfy condition (iv). On the other hand, $\alpha_{s_1+1} = k_2$ generates a root, therefore it has to be

$$\max\{\,\alpha_j\,|\,j \le s_1\,\} < k_2 - 1$$

Similarly, the second part, $(k_2, \ldots, \alpha_s)$ must not generate another root except in k_2 and therefore it has to satisfy (iv) as well. Consequently, the partition function for two roots has the form

$$\omega_n(\{k_1,k_2\},s) = \sum_{s_1+s_2=s} \omega_{k_2-2}(\{k_1\},s_1)\ \omega_n(\{k_2\},s_2) \tag{19}$$

The generating function reads

$$\Omega_n(\{k_1,k_2\},\lambda) = \Omega_{k_2-2}(\{k_1\},\lambda)\ \Omega_n(\{k_2\},\lambda) \tag{20}$$

In this way, we can proceed to heaps with arbitrarily many roots. The generating function of heaps with ℓ roots $k_1 < k_2 < \cdots < k_\ell$ reads

$$\Omega_n(\{k_1,k_2,\ldots,k_\ell\},\lambda) = \Omega_n(\{k_\ell\},\lambda) \prod_{i=1}^{\ell-1} \Omega_{k_{i+1}-2}(\{k_i\},\lambda) \tag{21}$$

Inserting (17), and setting

$$\begin{aligned} \Delta_0 &= k_1 && \text{(22a)}\\ \Delta_i &= k_{i+1} - k_i - 1 \quad \text{for} \quad 1 \le i < l && \text{(22b)}\\ \Delta_\ell &= n - k_\ell + 1 && \text{(22c)} \end{aligned}$$

we obtain

$$\Omega_n(\{k_1,\ldots,k_\ell\},\lambda) = \frac{1}{\sin(n+2)\phi} \prod_{i=0}^{\ell} \frac{\sin\Delta_i\phi}{\sin\phi} \tag{23}$$

for $\lambda \in [-1,3]$. For $\ell = 1$, equation (23) reproduces equation (17).

The final step is to apply the inverse Z–transform to the generating function (23) in order to obtain the partition function. This is done by performing the integral

$$\omega_n(C,s) = \frac{1}{2\pi i} \oint_{|\lambda|=\rho} \frac{\Omega_n(C,\lambda)}{\lambda^{-s+1}}\, d\lambda \tag{24}$$

where the contour $|\lambda| = \rho$ surrounds all poles

$$\lambda_m = 4\cos^2\frac{m\pi}{n+2} - 1 \quad ; \quad m = 1\dots\left\lfloor\frac{n+1}{2}\right\rfloor \tag{25}$$

of $\Omega_n(C,\lambda)$. Applying the residue theorem, we obtain

$$\omega_n(C,s) = \sum_{\lambda_m}\operatorname*{Res}_{\lambda_m}\left(\frac{1}{\sin(n+2)\phi}\prod_{i=0}^{\ell}\frac{\sin\Delta_i\phi}{\sin\phi}\right) \tag{26}$$

The residues can be easily evaluated, and the partition function of heaps on n columns with set of roots $C = \{k_1,\dots,k_\ell\}$ is derived as

$$\omega_n(C,s) = \sum_{m=1}^{2n+3}\frac{(-1)^{m+1}}{n+2}\sin\frac{2m\pi}{n+2}\left(\sin\frac{m\pi}{n+2}\right)^{-\ell}\prod_{i=0}^{\ell}\left(\sin\frac{\Delta_i m\pi}{n+2}\right)\left(1+2\cos\frac{2m\pi}{n+2}\right)^{s-1} \tag{27}$$

Combining (27) and (2), we arrive at our final result (1). It coincides with the one given in [2].

VI. SUMMARY

The number of directed animals with s occupied sites on a strip of width n for given root C has been calculated by using the analogy to the model of heaps of pieces. The result, which is formulated in theorem 1, coincides with the one obtained by Hakim and Nadal [2] who used an algebraic method dealing with the diagonalization of the transfer matrix for some spin system.

It has been shown by D. Dhar in [1] that the statistical weight of a configuration of two–dimensional directed animals is the same as the statistical weight of a spin configuration of an antiferromagnetic Ising model with anisotropic nearest–neighbor coupling in an external magnetic field on a triangular lattice with a Hamiltonian

$$H = \sum_{i,j}\left(L n_{i,j} n_{i-1,j+1} - \mu n_{i,j}\right) \tag{28}$$

The correspondence between directed animals and the 2D Ising spin system is set as follows. Define the fugacity, z, conjugated to the animal's size n (z is the same as λ^{-1} in our computations). Now the generating function, $A(z)$ of directed–site lattice animals on the square lattice reads $A(z) = -\partial_\alpha f(\alpha,z)|_{\alpha=0}$ where $\ln(1+z) + f(\alpha,z)$ is the Gibbs free energy per site for the antiferromagnetic Ising system with the Hamiltonian H defined above. The parameters α and z are connected to the coupling constant L and to the chemical potential μ in (28) by the equations $e^{-L} = 1+z$, $e^{\mu} = -(1-\alpha)z(1+z)^3$.

Note that $A(z)$ coincides up to the redefinition $z \to \lambda^{-1}$ with $\Theta_n(n,\lambda)$ defined in (10)–(11). The minor difference between $A(z)$ and $\Theta_n(n,\lambda)$ consists in the fact that $A(z)$ is defined for periodic boundary conditions, while $\Theta_n(n,\lambda)$ – for open ones.

By using the concept of the locally free group, the enumeration of directed lattice animals becomes a rather simple combinatoric problem. So, it seems an intriguing question how far the ideas of geometric group theory could be extended in the area of two–dimensional exactly solvable spin systems.

The proposed combinatorial approach to the statistics of lattice animals admits few possible generalizations. The simplest one consists in relaxing the idempotent condition $f_i^2 = f_i$ (see 3b), or replacing it by more general ones: either $f_i^p = f_i$ or $f_i^p = 1$ ($p = 1,2,\dots$). Such a modification of the problem allows to consider directed lattice animals on a class of decorated lattices. For example, skipping the relation $f_i^2 = f_i$, we pass to the triangular lattice – see Fig. 6a.



Figure 6: (a) Animals on triangular lattice; (b) Left–, Right– and Double–handed supports of particles on a square lattice.

A more involved generalization of the model allows to distinguish between "left–handed" (L) and "right–handed" (R) supports of deposited particles as shown in Fig. 6b. To proceed with the description, suppose that at the initial time $t = 0$ the system is empty. Then, at each time step, $t = 1,2,\dots,N$, an elementary unit ("particle") is deposited at some column. Consider the deposited particles as distinguishable (i.e. "colored"), and assign to them the variable t according to the natural time ordering. Now each particle carries two indices: the index t fixes its occurrence in time (i.e. a "color"), and the index α_i labels the column in which the particle is deposited. Now proceed with normal ordering, but apply it twice to the sequence: first – exactly as it is described in Definition 2, i.e. pushing the smaller indices α_i as left as possible (allowed by the commutation relations) and second – inversely, pushing the smaller indices α_i as right as possible (also consistent with the commutation relations). The first ordering will catch the L–supported particles, while the second ordering – the R–supported particles. Since the particles are colored, we can easily identify whether a given particle has L, R, or double supports. Introducing different weights $w_{\rm L}, w_{\rm R}, w_{\rm D}$ for L–, R– and double supports, we could distinguish between these configurations, which is crucial for a wide class of models (for example, for cellular automata [6]).

The authors are grateful to A. Vershik and R. Bikbov for helpful discussions. N.H. acknowledges the LPTMS, S.N is partially supported by the ANR grant 2011-BS04-013-01 “WALKMAT”.

---

[1] Dhar, D., Equivalence of the Two-Dimensional Directed-Site Animal Problem to Baxter's Hard-Square Lattice-Gas Model, Phys. Rev. Lett. **49** 959 (1982)

[2] Hakim, V.; Nadal, J.P., Exact results for 2D directed lattice animals on a strip of finite width, J. Phys. A: Math. Gen. **16** L213 (1983)

[3] Nechaev, S.; Voituriez, R., Random walks on three-strand braids and on related hyperbolic groups, J. Phys. A: Math. Gen. **36** 43 (2003)

[4] Vershik, A.M.; Nechaev, S.; Bikbov, R., Statistical Properties of Locally Free Groups with Applications to Braid Groups, Comm. Math. Phys. **212** 449 (2001)

[5] Viennot, G.X., eds. Labelle, G.; Leroux, P., Heaps of pieces I: Basic Definitions and Combinatorial Lemmas, Lecture Notes in Maths, n° **1234** 321 (1986)

[6] P. Rujàn, Cellular automata and statistical mechanical models, J. Stat. Phys., **49** 139 (1987)

[7] The labelling has to be done in such a way that the node with index 1 is connected to only one node in the second row.

[8] We do not distinguish between the actual generating function, which is defined as a series, and its maximum analytic continuation.